

will be carried out towards forming a conceptual approach to developing effective standardized protocols for providing rehabilitation assistance at different stages and periods of rehabilitation to those in need, including military personnel, taking into account modern international standards and strategies for providing rehabilitation assistance.

Key words: combat injury, servicemen, medical and psychological rehabilitation.

ORCID and contributionship / ORCID кожного автора та їх внесок до статті:

Zabolotna I. B.: [0000-0002-0159-2058](https://orcid.org/0000-0002-0159-2058)^{ABD}

Gushcha S. G.: [0000-0003-3097-5258](https://orcid.org/0000-0003-3097-5258)^{BCD}

Balashova I. V.: [0000-0002-7529-4045](https://orcid.org/0000-0002-7529-4045)^{BDE}

Bezverkhniuk T. V.: [0000-0002-2567-8729](https://orcid.org/0000-0002-2567-8729)^{AEF}

Conflict of interest / Конфлікт інтересів:

The authors declare that there is no conflict of interest in this article. / Автори заявляють про відсутність конфлікту інтересів в даній статті.

Corresponding author / Адреса для кореспонденції

Gushcha Serhiy Hennadiyovych / Гуща Сергій Геннадійович

SI «Ukrainian Scientific-Research Institute of Medical Rehabilitation and Balneology, the Ministry of Public Health of Ukraine» / ДУ «Український науково-дослідний інститут медичної реабілітації та курортології МОЗ України»
Ukraine, 65000, Odessa, 6 Lermontovsky 6 ave. / Адреса: Україна, 65014, м. Одеса, пров. Лермонтовський 6
Tel.: 0956262722 / Тел.: 0956262722

E-mail: gushchasergey11@gmail.com

A – Work concept and design, B – Data collection and analysis, C – Responsibility for statistical analysis, D – Writing the article, E – Critical review, F – Final approval of the article. / A – концепція роботи та дизайн, B – збір та аналіз даних, C – відповідальність за статичний аналіз, D – написання статті, E – критичний огляд, F – остаточне затвердження статті.

Received 21.11.2022 / Стаття надійшла 21.11.2022 року
Accepted 03.05.2023 / Стаття прийнята до друку 03.05.2023 року

DOI 10.29254/2077-4214-2023-2-169-74-81

UDC 616.832.006.311.013:616.018.007

Zaychenko G. V., Gorchakova N. O., Klymenko O. V., Shumeiko O. V., Babak V. V.

PHYSICOCHEMICAL, BIOCHEMICAL, PHARMACOLOGICAL PROPERTIES OF MAGNESIUM

Bogomolets National Medical University (Kyiv, Ukraine)

gorchakovan1941@gmail.com

The article discusses the role of magnesium in the body, which is one of the main intracellular elements that plays a leading role in the activity of vital organs, participates in many cellular functions including signal transmission, energy production, protein metabolism, and acts as a molecular stabilizer for RNA and DNA ribosomes. Hypomagnesemia leads to disturbances in the cardiovascular, nervous, immune, and digestive systems. Based on magnesium, mono (magnesium sulfate, magnesium citrate, etc.) and complex preparations (MagneB6, Maalox, ATP forte, etc.) have been created and are being created. Imbalances in magnesium content are associated with the development of cardiovascular diseases such as arterial hypertension, arrhythmias, heart failure, as well as the occurrence of neurodegenerative diseases of the CNS (parkinsonism, Alzheimer's disease, dementia), as magnesium is involved not only in nucleic acid metabolism but also in signal transmission in the CNS. Therefore, monitoring magnesium levels in the cardiovascular, nervous, and other systems may be a target for the influence of magnesium preparations. In individuals with digestive tract diseases, changes in magnesium content have also been identified in the cardiovascular system and CNS. Changes in magnesium content are associated with the occurrence of cardiovascular and digestive tract diseases in children, as well as the course of pregnancy in women. This indicates the need for further research into the properties of this trace element in normal and pathological conditions, as well as the search for new methodological approaches to determining its functional impact on the human body.

Key words: magnesium, physicochemical, biochemical, pharmacological properties.

Connection of the publication with planned research works.

The article is presented as part of the planned research work of the Department of Pharmacology on "Experimental Substantiation of Combined Use of Cardiotropic Drugs" (Registration No. 0111U009417).

Introduction.

In addition to vitamins and hormones, micro – and macroelements are necessary for the functioning of the

body and normal functioning of organs and systems. In previous publications, the pharmacological properties of zinc, iron, selenium, amino acids were revealed, and their role in the functioning of the body and the development of new drugs was determined.

At the same time, significant attention is currently focused on the role of magnesium, which is a key microelement in the body and is second only to potassium in terms of content. Magnesium, the twelfth element of the

periodic table, was first obtained in 1809 by English scientist Humphry Davy through electrolysis and was named "magnesium" from the French word "magnifique," which means "magnificent." This element justifies this name, as evidenced by its involvement in improving endothelial function and its role in the metabolism of vital organs [1-2]. The total content of magnesium in the body of an average adult human is about 1000 mmol or 24 grams. Cells contain about 40-50% of the total magnesium content, while the rest is found in muscles. 96% of magnesium is intracellular. This element has become known not only due to the introduction of magnesium-containing drugs into medical practice, but also due to their wide application in various industries and agriculture [3]. The literature data presented in the review determine whether the pathogen exhibits consequences of diseases associated with changes in magnesium content in the body [4-5].

The aim of the study.

To analyze the physicochemical, biochemical, pharmacological, clinical properties of magnesium and its preparations, and the spectrum of their application in clinical practice.

Object and methods of research.

The research was conducted based on a literature review, including PubMed, Google Scholar, Scopus, on the topic of the study.

Results of the research and their discussion.

Magnesium is considered as biometal, along with potassium, calcium, zinc, cobalt, manganese, molybdenum, iron, and copper. Although the content of biometals in the body is small, they determine the coordination of processes of excitation, contraction, relaxation of muscles, transmission of impulses in the nervous and cardiovascular systems, and functioning of other vital systems [6]. Magnesium ions play an important role in the regulation of almost all organs and systems. Magnesium acts as a structural component of cells, serving as a trigger, allosteric effector, or modulator of a wide range of energy and plastic reactions.

Comparison with calcium, magnesium ions are more hydrated and have a pronounced ability to form coordination bonds with biomembrane elements, forming coordination ligands. As the concentration of magnesium increases, its ability to form complexes decreases, while for calcium, it increases. Inorganic salts and hydroxides of magnesium are well or moderately soluble in water and are strong electrolytes. Even though the four elements – potassium, calcium, sodium, and magnesium – play an important role in the body, magnesium ranks second in terms of content in organs and systems after potassium [7]. Magnesium not only competes with calcium, but also prevents the influx of sodium into cells. This may be related to the unidirectional effect of magnesium and potassium on the activity of $\text{Na}^+\text{-K}^+\text{-ATPase}$, as well as the magnesium-sodium antiporter. For example, the introduction of magnesium prevents cardiomyocytes from losing potassium and calcium, and from the influx of sodium. Magnesium participates in energy metabolism, specifically in the metabolism of adenosine nucleotides, the conversion of ADP to ATP with the participation of inorganic phosphorus, and the breakdown of ATP to ADP through complex formation with both adenosine nucleotides.

Magnesium promotes the conversion of adenosine monophosphate into cyclic adenosine monophosphate

by influencing adenylate cyclase. Approximately five hundred proteins are associated with magnesium ions through the carboxyl group, and magnesium serves as a cofactor in about 350 enzymes. Magnesium interacts with membrane-polarizing groups. It also forms complexes with glucose, with the stability of magnesium-glucose complexes being greater than that of calcium, which is attributed to electrostatic interactions. For coordination compounds, the entropy contribution is higher for magnesium-glucose complexes, while the enthalpy contribution is higher for calcium-glucose complexes. Magnesium complexes with glucosamine are more stable than calcium-glucosamine complexes. Magnesium participates in glycolysis, oxidative phosphorylation, and is required for the synthesis of RNA and DNA, as well as for the functioning of mitochondria. Magnesium enzymes are involved in protein synthesis, muscle contraction, nervous system function, glycemic control, ionic fluxes, especially calcium channels [8].

After magnesium intake through water and food, a person weighing 70 kg contains 25-28 grams of magnesium in various tissues: 50-53% in bones, 27-28% in muscles, 19-20% in other soft tissues, 0.5% in erythrocytes, and 0.3-0.4% in blood plasma. Like calcium, magnesium participates in enterosorption in the gastrointestinal tract, renal reabsorption and secretion, and exchange processes in various organs, forming magnesium depots (e.g., in bones) [9]. This contributes to maintaining a beneficial level of calcium in bones and supporting the crystalline lattice of tooth enamel [10]. Hypermagnesemia, a condition where the total serum magnesium level exceeds 1.2-1.3 mmol/L, is a less common phenomenon. It can occur in chronic kidney diseases, acute renal failure, excessive magnesium therapy, epilepsy, eclampsia, and prolonged stress. Adrenergic stimulation of lipolysis can increase magnesium deficiency by forming complexes of magnesium with fatty acids. Thromboxanes and endothelins, which are formed because of fatty acid metabolism, contribute to this condition.

Physical and emotional stresses increase the need for magnesium. Magnesium plays a significant role in the functioning of neurotransmitter systems. It acts as an agonist of glutamate receptors and participates in the activation of ATP-dependent potassium channels, blocking the influx of calcium through voltage-gated calcium channels. Magnesium is an agonist of GABA-A receptors and an antagonist of angiotensin II receptors, activating neurotransmission associated with protein kinase C function [11]. The use of lithium medications may lead to hypermagnesemia, where the level of magnesium in the blood increases parallel to the concentration of calcium.

Hypomagnesemia is more commonly encountered in clinical practice. Hypomagnesemia is observed in severe conditions of patients, including critical illnesses and certain diseases of vital systems and organs. This primarily applies to stress situations accompanied by increased levels of catecholamines and glucose. In this case, there is an increase in magnesium excretion in urine. Stress, elevated catecholamine levels, increase the frequency of cardiovascular events, including arterial hypertension. Hypomagnesemia is observed in cases of inadequate intake of magnesium from food, exposure to pesticides in the gastrointestinal tract, decreased vitamin D levels, consumption of hard water, and alcohol. Hypomagnesemia may be caused by medications such as H1 and H2 histamine

receptor blockers, beta 1 and alpha-adrenergic agonists, sodium bicarbonate, antibiotics, antituberculosis and antiviral drugs, methylxanthines, antiarrhythmic drugs, ACE inhibitors, cardiac glycosides, diuretics, CNS stimulants, glucocorticoids, immunosuppressants, progestins, estrogens, and proton channel blockers, androgens [12-14].

Magnesium is involved in cell proliferation processes, depending on the state of the nervous system and homeostasis. The magnesium content significantly decreases in severe diseases such as carcinogenesis and diabetes mellitus [2]. Studies on endothelial cells have shown that magnesium transports the TRPM7 potential receptor [15]. Magnesium content changes proportionally with changes in cognitive function [16]. The magnesium content in cerebrospinal fluid correlates with the magnesium content in erythrocytes, and the magnesium content in the pituitary gland determines the ability to memorize, which is somehow related to DNA structure. Studies focusing on determining magnesium levels in cells and intercellular fluid have established its direct influence on human health [17]. Magnesium deficiency is often associated with deficiencies in potassium, copper, zinc, and less commonly chromium, cobalt, or nickel in blood serum. It is known that a decrease in magnesium levels is accompanied by an increase in calcium concentration, which is observed in type 2 diabetes. It is important to note that magnesium deficiency contributes to potassium exchange disorders, which play a role in cardiovascular pathology. Decreased magnesium content may be associated with increased excitability of the nervous system, and in cardiomyocytes, it may lead to tachycardia and ectopic arrhythmia. Decreased magnesium concentration in smooth muscle cells leads to increased blood pressure and headaches, as well as cramps, abdominal pain, bronchospasm, and uterine hypertonicity [18]. Long-term magnesium deficiency can lead to the development of cardiovascular and cerebral pathological conditions, such as heart attack, stroke, and atherosclerosis. Magnesium deficiency not only contributes to the onset of these diseases, but also to their severe course. Currently, there are many magnesium preparations available in various doses and different pharmaceutical forms, including inorganic salts such as sulfate, chloride, hydroxide, oxide, and salts of organic acids. Inorganic magnesium salts have low bioavailability, which is why magnesium sulfate is prescribed as a laxative and detoxification agent. Organic magnesium salts have significantly higher bioavailability, making them more relevant for use, such as magnesium citrate, lactate, orotate, pyroglutamate, and others. Organic magnesium salts are better absorbed, better tolerated, and less likely to cause side effects. Treatment with magnesium will be more effective when combined with magnesium fixatives such as vitamins B1 and B6, glycine, orotic acid, and others. Different magnesium salts have their own indications, for example, magnesium citrate is used in nephrolithiasis, and magnesium hydroxide is used as an antacid.

Vitamin B6 is the main regulator of magnesium in the human body, as both magnesium and B6 are important for the prevention of endothelial dysfunction in arterial hypertension, carbohydrate and lipid metabolism, and mitochondrial function. There is a correlation between the development of ischemic heart disease and these micronutrients. Therefore, Magnesium-B6, a preparation consisting of magnesium citrate and pyridoxine, is used

to replenish magnesium stores [19]. Aspartate (Panangin), which contains potassium and magnesium aspartates, also has high bioavailability. Aspartate anions are absorbed by the body and enter metabolic pathways (amino acid metabolism, urea cycle, Krebs cycle) without burdening the excretory systems, and intensify the detoxification of nitrogen metabolism products in the kidneys. This is important for the prevention of arterial hypertension, as increased levels of uric acid and decreased magnesium levels can trigger arterial hypertension. Therefore, magnesium and potassium ions are necessary for reducing and normalizing systolic and diastolic blood pressure in hypertension. Systematic depletion of magnesium stores contributes to the development of chronic endothelial inflammation, increased risk of cardiovascular and cerebrovascular diseases, and may also lead to insulin resistance, intolerance, and diabetes [19-23].

In this pathology, magnesium deficiency is associated with intensified free radical reactions, oxidative stress, and inflammation. Magnesium preparations are effective in blocking IL-1 β , one of the main inflammatory factors. These preparations also have anti-inflammatory effects by suppressing the membrane cytoplasmic P2X7R receptor [24]. Increasing magnesium concentration in body fluids protects against oxidative stress and changes in immune competent cells [25]. In heart failure, plasma magnesium concentration is inversely correlated with the level of C-reactive protein and tumor necrosis factor (TNF α). Decreased plasma magnesium levels are associated with increased levels of monocyte chemoattractant protein-1 (MCP-1) and interleukins, such as IL-1, IL-6, TNF α , IL-8, MCP-1, in the blood.

In heart failure, several interconnected metabolic changes occur:

1. Induction of oxidative stress, as significant amounts of magnesium are consumed during stress [26].
2. Activation of the renin-angiotensin-aldosterone system.
3. Weakening of calcium channel inhibition.
4. Activation of phagocytosis.
5. Diminished antagonism of magnesium towards NMDA receptors, resulting in excitotoxicity and hyperproduction of pro-inflammatory substance P.
6. Activation of NF- κ B-mediated signaling and increased transcription of cytokines and pro-inflammatory genes.
7. Reduction in the levels of anti-inflammatory mediators such as NO, lipoxins, resolvins, and protectins.

Low magnesium levels increase the oxidative activity of neutrophils, while high magnesium levels significantly reduce the production of free radicals in rats and polymorphonuclear cells in humans [27].

Loss of magnesium content in the myocardium has been observed in myocardial infarction and acute heart failure in the area of necrosis formation. It has been noted that magnesium prevents the occurrence of arrhythmias, whereas magnesium deficiency can lead to fatal arrhythmias. Magnesium can strengthen the potential of ventricular myocardium, as low magnesium levels in cardiomyocytes lead to membrane destabilization, while high magnesium levels stabilize the membranes. Therefore, there is accumulating evidence that magnesium preparations can prevent arrhythmias. They are effective in various types of tachyarrhythmias, including those caused by digoxin, multifocal atrial tachycardia in-

duced by neuroleptics, as well as those observed before surgery, in emergency treatment of arrhythmias caused by antiarrhythmic drugs, and in the postoperative period.

Magnesium also has anti-ischemic effects. In patients with variant angina, a decrease in magnesium levels correlating with the development of vasospasm has been observed. A decrease in magnesium levels has also been documented in the diagnosis of chest pain and ST segment deviations during coronary spasm. There are studies that demonstrate the effectiveness of magnesium preparations in patients with myocardial infarction and ST segment elevation [28].

One of the well-known effects of magnesium is its ability to lower blood pressure. The mechanism of this effect is explained by its natural blockade of calcium channels. Magnesium also competes with sodium for binding sites on smooth muscle cells of blood vessels. Additionally, it increases the production of prostaglandin E through its one-way influence on potassium, leading to endothelium-dependent vasodilation and correction of endothelial dysfunction in patients with arterial hypertension and diabetes, thereby reducing calcium and sodium levels. Magnesium is also an important factor in the enzyme delta-6-desaturase, which limits the rate of conversion of linolenic acid to gamma-linolenic acid. Gamma-linolenic acid, in turn, elongates to dihomo-gamma-linolenic acid, which is a precursor of prostaglandin E1 (PGE1) that has vasodilatory effects and inhibits platelet aggregation [29-30].

Magnesium preparations are effective as antiarrhythmic agents. Magnesium orotate, like other magnesium preparations, has an antiarrhythmic effect, as well as membrane-stabilizing properties. Magnesium participates in the depolarization of cardiomyocytes and maintains electrolyte balance. The early magnesium preparations (sulfate, oxide) had an absorption rate of about 5%, but with the introduction of lactate and citrate, absorption increased to 50%. Organic acid salts (aspartate, orotate, glycine) were well absorbed without a laxative effect. Magnesium orotate prevented and alleviated ventricular arrhythmia in experiments on rats [31-32]. The preparation modulated the duration of action potential and reduced myocardial excitability, increased PR and QRS intervals, prolonged the refractory period of atria and ventricles, thus reducing arrhythmogenic effects. This effect was mediated through the influence on T and L calcium channels, as well as Na⁺-K⁺-ATPase, which prevented excessive potassium expression.

A deficiency of magnesium leads to decreased platelet aggregation. Additionally, a decrease in magnesium content worsens the outcome of hypokalemia, as magnesium deficiency disrupts the function of Na⁺-K⁺-ATPase, leading to reduced uptake of potassium by cells and decreased potassium content. The use of beta 1-adrenergic blockers, as well as disturbances in magnesium intake or expression, may contribute to the development of hypomagnesemia. Magnesium deficiency can also result in cardiomyopathy. In patients with cardiomyopathy, both in animals and humans, magnesium deficiency has been observed, accompanied by a dose-dependent reduction in the size of the lesion. Studies have shown that magnesium not only reduces the risk of developing cardiomyopathy and endocardial fibrosis, but also maintains ATP levels in the body. In addition, inadequate magnesium

content is associated with mitochondrial dysfunction [33].

Deficiency of magnesium has been shown to play a role in the survival of patients with cardiomyopathy and heart failure [32]. It has also been shown that patients with mitral valve prolapse have magnesium deficiency, which is associated with reduced magnesium content in lymphocytes, leading to induction of fibrosis and depletion of glycogen stores.

Magnesium is a natural anti-stress factor that inhibits the development of excitatory processes in the central nervous system (CNS) and reduces the sensitivity of the body to external influences. While calcium and glutamate are excitatory substances and can be toxic in excess, magnesium protects NMDA receptors from toxin action and ensures activation of the neuroprotective agent glycine [34]. The recommended daily intake of magnesium is 400 mg for men and 500 mg for women. Magnesium deficiency is more common in women, and the need for magnesium increases during pregnancy and lactation. Special attention should be paid to magnesium supplementation in pregnant women, as this mineral is essential for stabilizing the processes of mitosis and meiosis, which contribute to the formation of a healthy embryo. Magnesium is found in significant amounts in the heart, brain, muscles, but its highest concentration is in the placenta [33]. Therefore, magnesium deficiency in pregnant women results in placental pathology, inhibition of physiological apoptosis leading to embryonic developmental defects, alkaline phosphatase activity alteration in the placenta negatively affecting phosphate metabolism and transport of class G globulins. In addition, pregnant women with magnesium deficiency may experience seizures, more commonly in calf muscles, arrhythmias, and increased excitability. Moreover, it has been observed that magnesium deficiency in pregnancy leads to an increase in the level of cascade apoptosis [34]. At the same time, mice that received magnesium acetyl taurate at all doses showed an increase in magnesium levels in brain tissue and skeletal muscles. Moreover, the increase in magnesium levels was more pronounced after a single administration of the preparation compared to twice-daily administration. The mechanisms of magnesium transport were found to be dependent on electrochemical gradient, transmembrane transport, and the soluble form of magnesium. Symptoms of magnesium deficiency and stress are very similar and are accompanied by dizziness, anxiety, uncontrollable weakness, and headache. Magnesium penetrates very well through the blood-brain barrier, controls membrane excitability, and is present in significant amounts in the extracellular space and cerebrospinal fluid, playing a significant role in brain homeostasis. In cerebrospinal fluid, magnesium exists in free and protein-bound forms [34]. Hypomagnesemia carries not only the risk of neurological and psychiatric disorders, but also type 2 diabetes, metabolic syndrome, osteoporosis, and cardiovascular diseases [2]. It has been established that magnesium deficiency not only leads to stress, but also to depression, cognitive memory disorders, and degenerative diseases. The manifestations of central nervous system disorders correlate with the duration of magnesium deficiency – prolonged magnesium deficiency leads to impaired hippocampal function and neurodegenerative and cognitive disorders [35]. Magnesium deficiency can also trigger bipolar disorders and

may require magnesium supplementation, especially in combination with vitamin B6 [36-38].

Research has shown that magnesium ions block NMDA-associated channels in a voltage-dependent manner and, by engaging in non-competitive antagonism with glutamate, inhibit its release, thereby reducing excitotoxicity [38]. Hypomagnesemia leads to genetic disorders of the central nervous system (CNS) that require magnesium therapy from childhood [39-40]. The neuroprotective effects of magnesium are associated with its role in genome repair and interaction with phosphate groups of DNA [41]. Magnesium deficiency can damage dopamine receptors and alter the sensitivity of NMDA and AMPA receptors [42]. Magnesium deficiency contributes to the development of Parkinson's disease, Alzheimer's disease, and dementia [43-44]. Genetic defects in the synthesis of dopamine transporters due to magnesium deficiency (TRPM7 and SCLA1) promote the development of Parkinson's disease [43-44]. In the pathogenesis of Alzheimer's disease, it has been noted that magnesium may prevent the formation of beta-amyloid and influence the NKB factor [43-44].

Magnesium relaxes smooth and skeletal muscles by inhibiting calcium-dependent channels, leading to the relaxation of myocytes. A comparison was made between the absorption of inorganic magnesium salts (magnesium citrate, magnesium malate) and organic salts (magnesium acetyl taurate, magnesium glycinate), with attention to magnesium transport, bioavailability, and impact on the immune system [42]. All compounds were administered in three different doses: 45 mg, 135 mg, and 405 mg of elemental magnesium per 70 kg of body weight. The highest dose was divided into 202.5 mg/70 kg and taken twice a day.

It is known that magnesium deficiency can occur due to inadequate intake from food, improper food preparation, certain diseases, or the use of certain medications. Magnesium galenic preparations were prepared and administered to mice, and magnesium levels were determined in brain, muscle tissues, and serum after one day [42]. Magnesium, along with vitamin supplements B, C, D, E, omega-3 polyunsaturated fatty acids, and manganese, may boost immunity. The mechanism of action of magnesium may be effective in patients with mutations in interleukin 2 and inducible forms of nitric oxide, as magnesium controls membrane excitability and can influence inflammation and immunity processes associated with these conditions. Therefore, it is considered that magnesium, along with the aforementioned vitamin supplements, can be prescribed in the treatment of viruses, infections, including influenza and COVID-19 [45-47]. Due to the association between psychoemotional and endocrine disorders with magnesium deficiency in the body, children with signs of chronic gastritis and arterial hypertension were additionally prescribed magnesium lactate with vitamin B6 according to the protocol, based on existing hypomagnesemia. This resulted in a reduction of abdominal and headaches, alleviation of stomach discomfort, improvement in sleep quality, and a decrease in daily systolic arterial tension (SAT) over the course of one month [48].

Magnesium plays a significant role in the detoxification process by promoting the elimination of toxic substances from the body. It is known that magnesium is necessary for the elimination of mercury, lead, zinc, and

arsenic from the body. Magnesium protects the body from the toxic effects of calcium, lead, mercury, nickel, and prevents the penetration of toxic metals into brain cells [49-51].

It is known that type 2 diabetes is a risk factor for the development of heart failure. In patients with type 2 diabetes, metabolic syndrome, and a history of ischemic heart disease, who had low levels of magnesium and zinc in their blood, zinc and magnesium supplements were added to the standard therapy. After receiving traditional therapy along with zinc and magnesium supplements, the overall condition of the patients improved, and the levels of magnesium, zinc, C-reactive proteins, LDL cholesterol, and markers of prooxidant-antioxidant homeostasis normalized. The obtained results indicated the effectiveness of the prescribed therapy [51-52].

Thus, magnesium actively regulates the state of the cellular membrane, facilitates the transfer of calcium and sodium ions, and is involved in metabolic reactions, including energy production. 80-90% of magnesium is complexed with ATP, which facilitates its accumulation in the body. Magnesium is necessary for nerve-muscle conduction, reduces anxiety, inhibits the release of catecholamines and aldosterone during stress states. Additionally, magnesium promotes reactions such as DNA synthesis, glycolysis, oxidative phosphorylation, as it is a component of guanosine triphosphatase, phosphofructokinase, Na-K-ATPase, Ca-ATPase, adenylate cyclase, and plays an important role in the transmission of genetic information, as it participates in nucleotide production [53]. Magnesium is crucial for the functioning of the cardiovascular system as it maintains myocardial function and exerts a hypotensive effect. Balanced magnesium levels are necessary for the functioning of the immune system and the support of B and T lymphocyte functions [53-55]. Considering the influence of magnesium on the immune system, magnesium supplements are recommended to be included in the treatment regimens of bronchial asthma in children [56].

In recent years, it has been determined that magnesium preparations can have an antinociceptive effect, which is associated with their influence on NMDA receptors, specifically inhibiting their activity. This allows for the use of these preparations in children and pregnant women [57, 58].

The data provided indicate that there is no system in the body that does not involve the participation of magnesium ions in the functioning of metabolic and genomic processes of organs and systems.

Conclusions.

Thus, magnesium is one of the key intracellular elements that plays a leading role in the activity of vital organs, participates in many cellular functions, including signal transmission, energy production, protein metabolism, and acts as a molecular stabilizer for RNA and DNA ribosomes. Hypomagnesemia leads to disruptions in the functioning of the cardiovascular, nervous, and immune systems, as well as the gastrointestinal tract. Based on magnesium, mono-preparations (such as magnesium sulfate, magnesium citrate, and others) and complex preparations (such as MagneB6, Maalox, ATP forte, and others) have been developed and continue to be developed. The development of cardiovascular diseases such as arterial hypertension, arrhythmias, heart failure, as well as neurodegenerative diseases of the central nervous system

(Parkinsonism, Alzheimer's disease, dementia) is associated with disturbances in magnesium levels, as magnesium participates not only in nucleic acid metabolism but also in signal transmission in the central nervous system. Therefore, monitoring magnesium levels not only in the cardiovascular and nervous systems but also in other systems may be a target for the effects of magnesium preparations. In individuals with gastrointestinal disorders, disturbances in magnesium levels have been identified not only in the gastrointestinal tract but also in the cardiovascular system and central nervous system. Low

magnesium levels are associated with the occurrence of cardiovascular and gastrointestinal diseases in children, as well as pregnancy outcomes in women. This indicates the necessity for further research into the properties of this micronutrient in health and disease, and the search for new methodological approaches to determine its impact on the human body.

Prospects for further research.

The role of metabolite-based drugs in the development of modern pharmaceuticals will be further investigated.

References

- Cunha AR, D'El-Rei J, Medeiros F, Umbelino B, Oigman W, Touyz RM, et al. Oral magnesium supplementation improves endothelial function and attenuates subclinical atherosclerosis in thiazide-treated hypertensive women. *Journal of Hypertension*. 2017;35(1):89-97. DOI: [10.1097/hjh.0000000000001129](https://doi.org/10.1097/hjh.0000000000001129).
- de Baaij JH, Hoenderop JG, Bindels RJ. Magnesium in man: Implications for health and disease. *Physiological Reviews*. 2015;95(1):1-46. DOI: [10.1152/physrev.00012.2014](https://doi.org/10.1152/physrev.00012.2014).
- Kursov SV, Nikonov VV, Biletskyi OV, Fedets OI, Homenko VO. Fiziologiya obmena magniyai primenenie magnezii v intensivnoj terapii (literaturnyj obzor s rezultatsami sobstvennykh nablyudenij, chast 1). *Medytsyna nevidkladnykh staniv*. 2022;17(5):56-67. DOI: [10.22141/2224-0586.17.5.2021.240708](https://doi.org/10.22141/2224-0586.17.5.2021.240708).
- Nosenko OM, Khancha FO. Mahnii u profilaktytsi hestatsiinykh uskladnenu zhinko iz vahitnistiu, indukovanioiu v tsykli DRT. *Reproduktyvna endokrynolohia*. 2020;(56):80-87. DOI: [10.18370/2309-4117.2020.56.80-87](https://doi.org/10.18370/2309-4117.2020.56.80-87). [in Ukrainian].
- Viering DH, de Baaij JH, Walsh SB, Kleta R, Bockenhauer D. Genetic causes of hypomagnesemia, a clinical overview. *Pediatric Nephrology*. 2017;32(7):1123-1135. DOI: [10.1007/s00467-016-3416-3](https://doi.org/10.1007/s00467-016-3416-3).
- Chekman IS, Gorchakova NA, Nikolay SL. Magnij v medicine. Kishinev; 1992. 101 s.
- Fiorentini D, Cappadone C, Farruggia G, Prata C. Magnesium: Biochemistry, nutrition, detection, and social impact of diseases linked to its deficiency. *Nutrients*. 2021;13(4):1136. DOI: [10.3390/nu13041136](https://doi.org/10.3390/nu13041136).
- Biletskyi OV. Zmina kontsentratsii elektrolitiv plazmy krovi v umovakh mahnezialnoi terapii u patsientiv u stani travmatychnoho shoku na tli politravmy. *Medytsyna nevidkladnykh staniv*. 2021;(5.100):69-73. DOI: [10.22141/2224-0586.5.100.2019.177021](https://doi.org/10.22141/2224-0586.5.100.2019.177021). [in Ukrainian].
- Al Alawi AM, Majoni SW, Falhammar H. Magnesium and human health: Perspectives and Research Directions. *International Journal of Endocrinology*. 2018;2018:1-17. DOI: [10.1155/2018/9041694](https://doi.org/10.1155/2018/9041694).
- Klitynska OV, Stishkovskyy AV. Mahnii v orhanizmi ta yoho rol u formuvanni stomatolohichnoi zakhvoriuvanosti. *Ukrayina. Zdorovya natsiyi*. 2020;3(60):130-137. DOI: [10.24144/2077-6594.3.2020.208661](https://doi.org/10.24144/2077-6594.3.2020.208661). [in Ukrainian].
- Chiarello DI, Marin R, Proverbio F, Benzo Z, Piñero S, Botana D, et al. Effect of hypoxia on the calcium and magnesium content, lipid peroxidation level, and Ca²⁺-ATPase activity of syncytiotrophoblast plasma membranes from placental explants. *BioMed Research International*. 2014;2014:1-9. DOI: [10.1155/2014/597357](https://doi.org/10.1155/2014/597357).
- Gröber U. Magnesium and drugs. *International Journal of Molecular Sciences*. 2019;20(9):2094. DOI: [10.3390/ijms20092094](https://doi.org/10.3390/ijms20092094).
- Nielsen FH. Magnesium deficiency and increased inflammation: Current perspectives. *Journal of Inflammation Research*. 2018;11:25-34. DOI: [10.2147/jir.s136742](https://doi.org/10.2147/jir.s136742).
- Yamanaka R, Shindo Y, Oka K. Magnesium is a key player in neuronal maturation and Neuropathology. *International Journal of Molecular Sciences*. 2019;20(14):3439-3465. DOI: [10.3390/ijms20143439](https://doi.org/10.3390/ijms20143439).
- Zhu D, Su Y, Fu B, Xu H. Magnesium reduces blood-brain barrier permeability and regulates amyloid- β transcytosis. *Molecular Neurobiology*. 2018;55(9):7118-7131. DOI: [10.1007/s12035-018-0896-0](https://doi.org/10.1007/s12035-018-0896-0).
- Sun Q, Weinger JG, Mao F, Liu G. Regulation of structural and functional synapse density by L-threonate through modulation of intraneuronal magnesium concentration. *Neuropharmacology*. 2016;108:426-439. DOI: [10.1016/j.neuropharm.2016.05.006](https://doi.org/10.1016/j.neuropharm.2016.05.006).
- Safaryan AS, Sargsyan VS, Kamyshova TV, Akhmedzhanov NM, Nebieridze DV, Poddubskaya EA. Rol magniya v razvitii serdechno-sosudistoj patologii i vozmozhnosti ee predotvrasheniya i korrektsii preparatami magniya. *Racionalnaya farmakoterapiya v kardiologii*. 2019;15(5):725-735. DOI: [10.20996/1819-6446-2019-15-5-725-735](https://doi.org/10.20996/1819-6446-2019-15-5-725-735).
- Zakharova IN, Tvorogova TM, Stepurina LL, Elezova LI. Rol defitsita magniya v formirovanii funktsionalnogo zapora u detej i podrostkov. *Medicinskij sovet*. 2019;(11):108-116. DOI: [10.21518/2079-701x-2019-11-108-116](https://doi.org/10.21518/2079-701x-2019-11-108-116).
- Gromova OA, Torshin IY, Nazarenko AG, Kalachev AGK. Deficit magniya i piridoksina kak faktory riska ishemicheskoy bolezni serdca. *Kardiologiya*. 2016;10:55-62. DOI: [10.18565/cardio.2016.10.55-62](https://doi.org/10.18565/cardio.2016.10.55-62).
- Yudina NV, Torshin IY, Gromova OA, Egorova EY, Bykov AT. Obespechennost ionami kaliya i magniya – fundamentalnoe uslovie dlya podderzhanija normalnogo arterialnogo davleniya. *Kardiologiya*. 2016;10:80-89. DOI: [10.18565/cardio.2016.10.80-89](https://doi.org/10.18565/cardio.2016.10.80-89).
- Sun H-L, Pei D, Lue K-H, Chen Y-L. Uric acid levels can predict metabolic syndrome and hypertension in adolescents: A 10-year longitudinal study. *PLOS ONE*. 2015;10(11):e0143786. DOI: [10.1371/journal.pone.0143786](https://doi.org/10.1371/journal.pone.0143786).
- Pilchova I, Klacanova K, Tatarkova Z, Kaplan P, Racay P. The involvement of Mg²⁺ in regulation of cellular and mitochondrial functions. *Oxidative Medicine and Cellular Longevity*. 2017;2017:1-8. DOI: [10.1155/2017/6797460](https://doi.org/10.1155/2017/6797460).
- Zhang G, Gruskos JJ, Afzal MS, Buccella D. Visualizing changes in mitochondrial Mg²⁺ during apoptosis with organelle-targeted triazole-based ratiometric fluorescent sensors. *Chemical Science*. 2015;6(12):6841-6846. DOI: [10.1039/c5sc02442k](https://doi.org/10.1039/c5sc02442k).
- Xie C, Li X, Zhu J, Wu J, Geng S, Zhong C. Magnesium isoglycyrrhizinate suppresses LPS-induced inflammation and oxidative stress through inhibiting NF- κ B and MAPK pathways in raw264.7 cells. *Bioorganic & Medicinal Chemistry*. 2019;27(3):516-524. DOI: [10.1016/j.bmc.2018.12.033](https://doi.org/10.1016/j.bmc.2018.12.033).
- Ozen M, Xie H, Shin N, Al Yousif G, Clemens J, McLane MW, et al. Magnesium sulfate inhibits inflammation through P2X7 receptors in human umbilical vein endothelial cells. *Pediatric Research*. 2019;87(3):463-471. DOI: [10.1038/s41390-019-0557-7](https://doi.org/10.1038/s41390-019-0557-7).
- Pickering G, Mazur A, Trousselard M, Bienkowski P, Yaltseva N, Amessou M, et al. Magnesium status and stress: The Vicious Circle Concept revisited. *Nutrients*. 2020;12(12):3672. DOI: [10.3390/nu12123672](https://doi.org/10.3390/nu12123672).
- Liu M, Dudley SC. Magnesium, oxidative stress, inflammation, and cardiovascular disease. *Antioxidants*. 2020;9(10):907. DOI: [10.3390/antiox9100907](https://doi.org/10.3390/antiox9100907).
- Szapary LB, Szakacs Z, Farkas N, Schonfeld K, Babocsay D, Gajer M, et al. The effect of magnesium on reperfusion arrhythmias in STEMI patients, treated with PPCI. A systematic review with a meta-analysis and trial sequential analysis. *Frontiers in Cardiovascular Medicine*. 2021;7:608193. DOI: [10.3389/fcvm.2020.608193](https://doi.org/10.3389/fcvm.2020.608193).
- Wongdee K, Rodrat M, Teerapornpuntakit J, Krishnamra N, Charoenphandhu N. Factors inhibiting intestinal calcium absorption: Hormones and luminal factors that prevent excessive calcium uptake. *The Journal of Physiological Sciences*. 2019;69(5):683-696. DOI: [10.1007/s12576-019-00688-3](https://doi.org/10.1007/s12576-019-00688-3).

30. Syrkin AL, Salagaev GI, Syrkina EA, Lysenko AV. Preimushstva orotata magniya dlya korrektsii magnij-defitsitnyh sostoyanij u bolnyh s razlichnymi formami narushenij ritma serdca. Kardiologiya i serdechno-sosudistaya khirurgiya. 2019;12(4):308. DOI: [10.17116/kardio201912041308](#).
31. Song EK, Kang S-M. Micronutrient deficiency independently predicts adverse health outcomes in patients with heart failure. Journal of Cardiovascular Nursing. 2017;32(1):47-53. DOI: [10.1097/jcn.0000000000000304](#).
32. Sontia B, Touyz RM. Role of magnesium in hypertension. Archives of Biochemistry and Biophysics. 2007;458(1):33-39. DOI: [10.1016/j.abb.2006.05.005](#).
33. Liu M, Liu H, Feng F, Xie A, Kang GJ, Zhao Y, et al. Magnesium deficiency causes a reversible, metabolic, diastolic cardiomyopathy. Journal of the American Heart Association. 2021;10(12):e020205. DOI: [10.1161/jaha.120.020205](#).
34. Dikke GB. Use of magnesium in obstetrics: Myths and facts. Obstetrics, Gynecology and Reproduction. 2017;11(3):59-68. DOI: [10.17749/2313-7347.2017.11.3.059-068](#).
35. Lo Piano F, Corsonello A, Corica F. Magnesium and elderly patient: the explored paths and the ones to be explored: a review. Magnesium Research. 2019;32(1):1-15. DOI: [10.1684/mrh.2019.0453](#).
36. Guidi J, Lucente M, Sonino N, Fava GA. Allostatic load and its impact on Health: A Systematic Review. Psychotherapy and Psychosomatics. 2020;90(1):11-27. DOI: [10.1159/000510696](#).
37. Pouteau E, Kabir-Ahmadi M, Noah L, Mazur A, Dye L, Hellhammer J, et al. Superiority of magnesium and vitamin B6 over magnesium alone on severe stress in healthy adults with low magnesemia: A randomized, single-blind clinical trial. PLOS ONE. 2018;13(12):e0208454. DOI: [10.1371/journal.pone.0208454](#).
38. Kursov SV, Nikonov VV, Biletskyi OV, Fedets OI, Homenko VO. Fiziologiya obmena magniyai primenenie magnezii v intensivnoj terapii (literaturnyj obzor s rezul'tatami sobstvennyh nablyudenij, chast 3). Medytsyna nevidkladnykh staniv. 2022;17(8):33-46. DOI: [10.22141/2224-0586.17.5.2021.240708](#).
39. Semenchenko SI. Porivnialnyi vplyv ademolu, amantadynu sulfatu, mahniiu sulfatu na nevrolohichni defitsyt ta mnestychni funktsii u shchuriv iz modelli cherepno-mozkovoї travmy. Bukovynskiy medychnyj visnyk. 2019;23(4(92)):108-115. DOI: [10.24061/2413-0737.xxiii.4.92.2019.97](#). [in Ukrainian]
40. Antwi-Boasiako C, Kusi-Mensah YA, Hayfron-Benjamin C, Aryee R, Dankwah GB, Kwawukume LA, et al. Total serum magnesium levels and calcium-to-magnesium ratio in sickle cell disease. Medicina. 2019;55(9):547. DOI: [10.3390/medicina55090547](#).
41. Ates M, Kizildag S, Yuksel O, Hosgorler F, Yuce Z, Guvendi G, et al. Dose-dependent absorption profile of different magnesium compounds. Biological Trace Element Research. 2019;192(2):244-251. DOI: [10.1007/s12011-019-01663-0](#).
42. Ohno N, Ikenaka K. Axonal and neuronal degeneration in myelin diseases. Neuroscience Research. 2019;139:48-57. DOI: [10.1016/j.neures.2018.08.013](#).
43. Hamada AM. Vitamins, omega-3, magnesium, manganese, and thyme can boost our immunity and protect against COVID-19. European Journal of Biological Research. 2020;10(4):271-295. DOI: [10.5281/zenodo.3990659](#).
44. William JH, Danziger J. Magnesium deficiency and proton-pump inhibitor use: A clinical review. The Journal of Clinical Pharmacology. 2015;56(6):660-668. DOI: [10.1002/jcph.672](#).
45. Reddy P, Edwards LR. Magnesium supplementation in vitamin D deficiency. American Journal of Therapeutics. 2019;26(1):e124-e132. DOI: [10.1097/mjt.0000000000000538](#).
46. Hutten TJ, Sikma MA, Stokwieler RH, Wesseling M, Hoefer IE, Tiel Groenestege WM. Ionized and not total magnesium as a discriminating biomarker for hypomagnesemia in continuous venovenous haemofiltration patients. Nephrology Dialysis Transplantation. 2021;36(4):742-743. DOI: [10.1093/ndt/gfaa330](#).
47. Marushko YV, Zlobynets AS, Hyschak TV, Komissarova OS. Rezultaty zastosuvannya kombinovanoho preparatu mahniiu i vitaminu B6 u ditei iz poiednanyim perebihom khronichnoho hastroduodenitu ta pervynnoi arterialnoi hipertenzii. ZDOROVE ZHENSINNY. 2020;2(148):70-74. [in Ukrainian].
48. Boujelben M, Abdennabi R, Elfki A. The protective effect of Mg on accumulation biomarkers and markers of Cd-induced oxidative stress in adult male Wistar rats. Clinical Case Reports, Research & Trials. 2018;3:4-15. DOI: [10.31872/2018/CCRR-100128](#).
49. Lim AK, Azraai M, Pham JH, Looi WF, Wirth D, Ng AS, et al. Severe tachycardia associated with psychotropic medications in psychiatric inpatients: A Study of Hospital Medical Emergency Team Activation. Journal of Clinical Medicine. 2021;10(7):1534-1545. DOI: [10.3390/jcm10071534](#).
50. Stoner SC. Management of serious cardiac adverse effects of antipsychotic medications. Mental Health Clinician. 2017;7(6):246-254. DOI: [10.9740/mhc.2017.11.246](#).
51. Baker WL. Treating arrhythmias with adjunctive magnesium: Identifying future research directions. European Heart Journal – Cardiovascular Pharmacotherapy. 2016;15:108-117. DOI: [10.1093/ehjcvp/pvw028](#).
52. Hamedifard Z, Farrokhi A, Reiner Z, Bahmani F, Taghizadeh M, Ghotbi M, et al. The effects of combined magnesium and zinc supplementation on metabolic status in patients with type 2 diabetes mellitus and coronary heart disease. Lipids Health Dis. 2020;19:112-121. DOI: [10.21203/rs.2.21470/v1](#).
53. Myrachko UV. Korektsiia nedostatnosti mahniiu u ditei ta pidlitkiv z astenichnym syndromom i pervynnoi arterialnoi hipertenziei. Ratsionalna farmakoterapia. 2016;3(40):35-48. [in Ukrainian].
54. Riazanova OD, Reznichenko HI. Vplyv defitsytu mahniiu na proiavy stresu v zhynok reproduktyvnoho viku, yaki khvori na nespetsychni vahynity. Zaporizkyi medychnyj zhurnal. 2022;24(5):580-585. DOI: [10.14739/2310-1210.2022.5.252513](#). [in Ukrainian].
55. Özdemir A, Doğruel D. Efficacy of magnesium sulfate treatment in children with acute asthma. Medical Principles and Practice. 2020;29(3):292-298. DOI: [10.1159/000506595](#).
56. Shin H-J, Na H-S, Do S-H. Magnesium and Pain. Nutrients. 2020;12(8):2184. DOI: [10.3390/nu12082184](#).
57. Wright RHG, Le Dily F, Beato M. ATP, Mg²⁺, nuclear phase separation, and Genome Accessibility. Trends in Biochemical Sciences. 2019;44(7):565-574. DOI: [10.1016/j.tibs.2019.03.001](#).
58. Marushko IV, Hyschak TV. Obosnovanie primeneniya preparatov magniya v pediatricheskoj praktike (obzor literatury). SOVREMENNAYA PEDIATRIYA. 2016;78(6):27-32. DOI: [10.15574/sp.2016.78.27](#).

ФІЗИКО-ХІМІЧНІ, БІОХІМІЧНІ, ФАРМАКОЛОГІЧНІ ВЛАСТИВОСТІ МАГНІЮ

Зайченко Г. В., Горчакова Н. О., Клименко О. В., Шумейко О. В., Бабак В. В.

Резюме. Магній є одним з головних внутрішньоклітинних елементів, який грає одну з провідних ролей в діяльності життєво важливих органів, бере участь в багатьох функціях клітин, включаючи передачу збудження, утворення енергії, обмін білків, є молекулярним стабілізатором РНК та ДНК рибосом. Гіпомagneмія веде до порушення діяльності серцево-судинної, нервової та імунної систем, органів травного каналу. На підставі магнію створені і створюються моно (магнію сульфат, магнію цитрат та інші) і комплексні препарати (МагнеВ6, Маалокс, АТФ форте та інші).

З порушенням вмісту магнію пов'язаний розвиток таких серцево-судинних захворювань як артеріальна гіпертензія, аритмії, серцева недостатність, а також виникнення нервово-дегенеративних хвороб ЦНС (паркінсонізму, хвороби Альцгеймера, розвитку деменції), адже магній бере участь не тільки в нуклеїновому обміні, але і передачі сигналів в ЦНС. Тому контроль рівня магнію не тільки в серцево-судинній, нервовій та

інших системах може бути мішенню впливу препаратів магнію. Магній захищає організм від токсичного впливу кальцію, свинця, ртуті, нікелю, а також запобігає проникненню в клітини мозку токсичних металів. У осіб, що страждають захворюваннями травного каналу, визначали порушення вмісту магнію не тільки в травному каналі, а також в серцево-судинній системі та в ЦНС. Зі зниженням вмістом магнію пов'язують виникнення захворювань серцево-судинної системи і травного каналу у дітей, а також перебіг вагітності у жінок. Магній також грає важливу роль у функціонуванні серцево-судинної системи: підтримує функцію міокарду, реалізує гіпотензивний ефект. Збалансований рівень магнію необхідний для функціонування імунної системи. Зважаючи на наукові дані щодо впливу магнію на імунну систему, рекомендовано включати препарати магнію в схеми лікування бронхіальної астми у дітей.

Наведені дані свідчать про те, що немає жодної системи організму, в функціонування якої не брали участь іони магнію, що забезпечують метаболічні та взагалі геномні процеси органів і систем. Це свідчить про необхідність подальшого дослідження властивостей цього мікроелементу в нормі і патології та пошук нових методичних підходів до визначення його впливу на організм людини.

Ключові слова: магній, фізико-хімічні, біохімічні, фармакологічні властивості.

PHYSICOCHEMICAL, BIOCHEMICAL, PHARMACOLOGICAL PROPERTIES OF MAGNESIUM

Zaychenko G. V., Gorchakova N. O., Klymenko O. V., Shumeiko O. V., Babak V. V.

Abstract. Magnesium is one of the main intracellular elements that plays a prominent role in the functioning of vital organs, participating in various cellular functions, including signal transmission, energy production, protein metabolism, and acting as a molecular stabilizer of RNA and DNA ribosomes. Hypomagnesemia can lead to disruptions in the functioning of the cardiovascular, nervous, and immune systems, as well as the digestive tract organs. Based on magnesium, mono preparations (such as magnesium sulfate, magnesium citrate, etc.) and complex drugs (Magnesium-B6, Maalox, ATP Forte, etc.) have been developed and are being used.

The development of cardiovascular diseases such as arterial hypertension, arrhythmias, heart failure, as well as neurodegenerative diseases of the central nervous system (Parkinsonism, Alzheimer's disease, dementia) is associated with magnesium deficiency, as magnesium not only participates in nucleic acid metabolism, but also in signal transmission in the central nervous system. Therefore, monitoring magnesium levels not only in the cardiovascular and nervous systems but also in other systems may be targeted for the influence of magnesium preparations. Magnesium protects the body from the toxic effects of calcium, lead, mercury, nickel, and prevents the penetration of toxic metals into brain cells. Individuals with gastrointestinal disorders have been found to have magnesium deficiency not only in the gastrointestinal tract but also in the cardiovascular system and CNS. Reduced magnesium levels have been associated with the occurrence of cardiovascular and gastrointestinal diseases in children, as well as pregnancy outcomes in women. Magnesium also plays a crucial role in the functioning of the cardiovascular system, maintaining myocardial function, and implementing a hypotensive effect. Balanced magnesium levels are necessary for the functioning of the immune system. Based on scientific data regarding the influence of magnesium on the immune system, the inclusion of magnesium preparations in the treatment regimens for childhood bronchial asthma is recommended.

The data presented indicate that there is no system in the body in which magnesium ions do not participate, as they are involved in metabolic and genomic processes of organs and systems in general. This highlights the need for further research into the properties of this trace element in normal and pathological conditions, and the search for new methodological approaches to determine its impact on the human body.

Key words: magnesium, physicochemical, biochemical, pharmacological properties.

ORCID and contributionship:

Zaychenko G. V.: [0000-0002-3506-4800](https://orcid.org/0000-0002-3506-4800) ^{ABCF}

Gorchakova N. O.: [0000-0001-7311-7347](https://orcid.org/0000-0001-7311-7347) ^{ABCDE}

Klymenko O. V.: [0000-0002-2537-7029](https://orcid.org/0000-0002-2537-7029) ^B

Shumeiko O. V.: [0000-0003-0655-0911](https://orcid.org/0000-0003-0655-0911) ^B

Babak V. V.: — ^B

Conflict of interest:

The authors of the paper confirm the absence of conflict of interest.

Corresponding author

Gorchakova Nadia Oleksandrivna
Bogomolets National Medical University
Ukraine, 01601, Kyiv, 34 Berestey Avenue
Tel.: +380677925337
E-mail: gorchakovan1941@gmail.com

A — Work concept and design, B — Data collection and analysis, C — Responsibility for statistical analysis, D — Writing the article, E — Critical review, F — Final approval of the article.

Received 20.11.2022

Accepted 05.05.2023